



Effect of Noni Fruit Extract (*Morinda citrifolia*) on Glucose Intake to Diabetes Mellitus White Rat Muscle Tissue

Patricia Wulandari^{1*}

¹ Cattleya Consultation Center, Palembang, Indonesia

ARTICLE INFO

Keywords:

Morinda
Alloxan
Diabetes mellitus
Metformin
Glucose transporter type 4

*Corresponding author:

Patricia Wulandari

E-mail address:

dr.patricia.wulandari@gmail.com

The author has reviewed and approved the final version of the manuscript.

<https://doi.org/10.37275/ehi.v1i1.2>

ABSTRACT

Noni fruit (*Morinda citrifolia*) is one of the most common plants in Indonesia. This plant is often found in various regions in Indonesia, where these plants are often in the form of shrubs or wild plants that grow in yards or plantations. This study aims to assess the effect of noni (*Morinda citrifolia*) fruit extract on blood sugar levels and the expression of GLUT4 protein in muscle tissue which shows the potential of the test extract's ability to improve blood glucose intake in cells so that it can maintain blood sugar regulation. This research was an experimental study that used white rats as research subjects. Induction of diabetes was done by injecting alloxan at a dose of 110 mg/kg BW intraperitoneally; then the white rats were given 10% glucose to drink. Alloxan-induced white rats showed a very significant increase in blood sugar levels, where the use of the drug metformin was able to reduce blood sugar levels significantly. In white rats induced with diabetes mellitus, there was a decrease in GLUT4 levels in muscle tissue. The administration of metformin drugs or noni fruit extracts shows the ability to increase levels of GLUT4 protein. In conclusion, noni fruit extract affects lowering blood sugar levels in diabetes mellitus white rats by increasing glucose intake in cells and tissues.

1. Introduction

Diabetes mellitus is a chronic condition that disturbs the body's blood sugar regulation. This disorder is characterized by a decrease in the ability of the body's cells to intake glucose into cells. Due to the failure of cells in glucose intake, glucose buildup occurs in the extracellular, namely in plasma. The buildup of glucose in the plasma causes an increase in blood glucose levels in the plasma so that it will interfere with blood flow, which led to blood viscosity and decreased blood flow to various cells and tissues. Impaired blood flow to various cells and tissues leads to a decrease in the supply of oxygen and nutrients to cells and tissues resulting in damage and cell death which leads to decreased tissue performance.¹⁻⁵

Noni fruit (*Morinda citrifolia*) is one of the most

common plants in Indonesia. This plant is often found in various regions in Indonesia, where these plants are often in the form of shrubs or wild plants that grow in yards or plantations. Noni fruit contains quite various secondary metabolites, and this plant is rich in flavonoids, alkaloids, terpenes, and glycosides. The content of these secondary metabolite compounds is believed to be rich in antioxidant effects so that it has the effect of being able to suppress various oxidative stress conditions that cause damage to various organs due to blood sugar dysregulation.⁶⁻⁹

This study aims to assess the effect of noni (*Morinda citrifolia*) fruit extract on blood sugar levels and the expression of GLUT4 protein in muscle tissue which shows the potential of the test extract's ability to improve blood glucose intake in cells so that it can

maintain blood sugar regulation.

2. Methods

Animal model

A total of 30 white rats (*Rattus norvegicus*) Wistar strains were obtained from the Eureka Research Laboratory (Palembang, Indonesia) weighing between 200 – and 250 grams. All experimental animals were kept in cages under controlled conditions of 12 hours of the light-dark cycle, temperature $22 \pm 1^\circ\text{C}$ and humidity 40-60%, and given ad libitum food. The research treatment and procedures have received approval from the medical research ethics committee of the Faculty of Medicine, Universitas Sriwijaya (No. 177/kptfkunsri- rsmh/2019).

Noni fruit extraction preparation

Noni fruit *Simplicia* was obtained from the Tangwangmangu Herbal Research Center, Karanganyar, Indonesia. The process of noni fruit extraction is carried out by maceration in which 500 grams of *Simplicia* are macerated with 96% ethanol for 72 hours. Next, do the separation between the pulp and the macerate. The macerate was then evaporated using a rotary evaporator (Shimadzu) to obtain a thick extract, noni fruit extract (EM).

Animal model diabetes melitus

After 1 week of adaptation, the mice were randomly divided into the following six groups, each containing 5 animals: Normal control group, diabetes group (negative control), diabetes + metformin group (Met; 45 mg / kg), Diabetes + EM (50 mg / kg), diabetes group + EM (100 mg / kg) and diabetes group + EM (200 mg / kg). Induction of diabetes was done by injecting alloxan at a dose of 110 mg/kg BW intraperitoneally; then the white rats were given 10% glucose to drink. The positive control group was treated with metformin (Dexa Medica, Indonesia) for 14 days. In the treatment group, noni fruit extract was carried out for 14 days. The mice were anesthetized by injecting 10% Chloral Hydrate (3.5 ml/kg) intraperitoneally. The rats were sacrificed by intraperitoneal injection of 10% chlorine

hydrate. Blood serum was taken through the orbital vein, and the femoral muscle was taken from the thigh of the white rat. The serum was then centrifuged at 10,000 rpm for 10 minutes, at 25°C , and the supernatant was stored at -20°C for analysis of blood sugar levels using the spectrophotometer method (Biorad). Meanwhile, the muscle tissue was evacuated, some of which were homogenized and centrifuged to obtain a supernatant and put it in a later RNA solution (Sigma Aldrich) and stored at -20°C , for ELISA examination of GLUT4 protein.

Enzyme-linked immunosorbent assays (ELISA) GLUT 4

GLUT4 levels in joint synovial fluid were checked with Rat ELISA GLUT4 (Cloud Clone), based on the protocol contained in the manufacturer's protocols. Briefly, 50 μl of standard diluent or serum samples were added to the well coated with anti-GLUT4 and incubated at 37°C for 30 minutes. After the plates were washed, 100 μl of the biotinylated antibody solution was added and incubated for 30 minutes at 37°C . After three washing, 50 μl of the avidin-peroxidase complex solution was added and incubated for 15 minutes at 37°C . After washing, 50 μl of tetramethylbenzidine color solution was added and incubated in the dark for 15 minutes at 37°C . Finally, 50 μl stop solution was added to stop the reaction, and the optical density (OD) was measured using an ELISA reader (Biorad), a wavelength of 450 nm.

Phytochemical test

The noni fruit extract was analyzed for phytochemical screening which included tannins, alkaloids, flavonoids, quinones, saponins, and steroids/triterpenoids. The noni fruit extract was separated using TLC as a stationary phase in the form of silica gel GF254 and a mobile phase in the form of n-hexane: chloroform: ethyl acetate (2: 5: 5).

Statistical analysis

All data were presented as mean \pm standard deviation, and all statistical analyzes were performed

with the SPSS 25 (IBM) program. One-way ANOVA followed by a post hoc analysis was carried out to assess the difference in mean expression levels of each protein. $P < 0.05$ was determined as an indication that there was a significant difference in mean levels.

3. Results and Discussion

Table 1 shows the potential of noni fruit extract on the blood sugar levels of white rats. Alloxan-induced white rats showed a very significant increase in blood sugar levels, where the use of the drug metformin was able to reduce blood sugar levels significantly even though they had not reached the target blood glucose target of less than 200 mg/dL. The treatment with noni fruit extract was able to reduce blood sugar levels significantly, were at the EM dose of 100 and 200 mg/kg BW it was able to reduce blood sugar levels to reach a target of below 200 mg/kg BW.

Table 2 shows the levels of GLUT4 in muscle tissue, where the GLUT4 protein is an essential transporter in the regulation of glucose intake into cells. Increased

expression of GLUT4 in a tissue indicates an increase in the ability of cells to intake glucose. In white rats induced with diabetes mellitus, there was a decrease in GLUT4 levels in muscle tissue. The administration of metformin drugs or noni fruit extracts shows the ability to increase levels of GLUT4 protein.¹⁰⁻¹³

Table 3 shows the secondary metabolite content of noni fruit extract. Noni fruit extract is rich in flavonoids. The dominant flavonoids in noni fruit extract are believed to be responsible for the effect of blood glucose regulation.

The content of flavonoids in the leading secondary metabolite compound is believed to play a role in blood glucose regulation. Flavonoids increase the expression of the GLUT 4 protein in muscle tissue. Increased expression of GLUT 4 causes an increase in glucose intake in cells. Increased glucose intake into cells, causes a decrease in the buildup of glucose outside the cells and interstitial tissue, which leads to a decrease in blood glucose levels.¹⁴⁻¹⁶

Table 1. Level of blood glucose in serum

No.	Group	Blood Glucose (mg/dL) \pm SD	P-Value*
1.	Control	96.86 \pm 8.41	0.00
2.	Diabetes	399.23 \pm 15.43	-
3.	Diabetes + Met	219.11 \pm 10.21	0.00
4.	Diabetes + EM 50	226.12 \pm 21.43	0.00
5.	Diabetes + EM 100	195.11 \pm 17.65	0.00
6.	Diabetes + EM 200	149.83 \pm 12.21	0.00

* VS Diabetes + Met; ANOVA. post hoc Bonferroni; $p < 0.05$

Table 2. Level of GLUT 4 in muscle

No.	Group	GLUT4 (pg/mL) \pm SD	P-Value*
1.	Control	228.26 \pm 3.41	0.00
2.	Diabetes	68.23 \pm 15.43	-
3.	Diabetes + Met	167.41 \pm 7.21	0.00
4.	Diabetes + ES 50	89.12 \pm 21.43	0.00
5.	Diabetes + ES 100	103.11 \pm 18.65	0.00
6.	Diabetes + ES 200	135.83 \pm 10.12	0.00

* VS Diabetes + Met; ANOVA, post hoc Bonferroni; $p < 0.05$

Table 3. Phytochemical test of noni fruit extract

Ingredients	Saponin	Alkaloid	Triterpenoid	Steroid	Flavonoid
EM	+	+	+	++	+++

4. Conclusion

Noni fruit extract affects lowering blood sugar levels in diabetes mellitus white rats by increasing glucose intake in cells and tissues.

5. References

1. Asgary S, Rahimi P, Mahzouni P, Madani H. Antidiabetic effect of hydroalcoholic extract of *Carthamus tinctorius* L in alloxan-induced diabetic rats. *J Res Med Sci.* 2012; 17: 386–392.
2. Dey L, Attele AS, Yuan CS. Alternative therapies for type 2 diabetes. *Altern Med Rev.* 2002; 7: 45–58.
3. Eddouks M, Lemhadri A, Michel JB. Hypolipidemic activity of aqueous extract of *Capparis spinosa* L in normal and diabetic rats. *J Ethnopharmacol.* 2005; 98: 345–50.
4. Eddouks M, Lemhardi A, Micel JB. Caraway and caper: potential antihyperglycaemic plants in diabetic rats. *J Ethnopharmacol.* 2004; 94: 143–148.
5. Fabiane K, Ricardo S, Oliveira T, Nagem TJ, Pinto AD, Oliveira MG, Soares JF. Effect of flavonoids morin; quercetin and nicotinic acid on lipid metabolism of rats experimentally fed with triton. *Braz Arch Biol Techn.* 2001; 44: 263–267.
6. Grover JK, Yadav S, Vats V. Medicinal plants of India with anti-diabetic potential. *J Ethnopharmacol.* 2002; 8: 81–100.
7. González-Villalpando C, López-Ridaura R, Campuzano JC, González-Villalpando ME. The status of diabetes care in Mexican population: Are we making a difference? Results of the National Health and Nutrition Survey 2006. *Salud Publica Mex.* 2010; 52: S36–46.
8. Inzuchi SE, Maggs DG, Spollett GR, Page SL, Rite FS, Walton V. Efficacy and metabolic effect of Metformin and troglitazon in type II diabetes mellitus. *N Engl J Med.* 1998; 338: 867–872.
9. Khan A, Anderson RA. Insulin potentiating factor (IPF) present in foods, species and natural products. *Pak J Nutr.* 2003; 2: 254–257. [Google Scholar]
10. Khanfar MA, Sabri SS, Zarga MH, Zeller KP. The chemical constituents of *Capparis spinosa* of Jordanian origin. *Nat Prod Res.* 2003; 17: 9–14.
11. Matsuyama T, Shoji K, Takase H, Kamimaki I, Tanaka Y, et al. Effects of phytosterols in diacylglycerol as part of diet therapy on hyperlipidemia in children. *Asia Pac J Clin Nutr.* 2007; 16: 40–48.
12. Matthaus B, Ozcan M. Glucosinolates and fatty acid, sterol, and tocopherol composition of seed oils from *Capparis spinosa* Var *spinosa* and *Capparis ovata* Desf. Var *canescens* (Coss.) Heywood. *J Agric Food Chem.* 2005; 53: 7136–7141.
13. Moon J1, Lee SM, Do HJ, Cho Y, Chung JH, et al. Quercetin up-regulates LDL Receptor Expression in HepG2 Cells. *Phytother Res.* 2012; 26: 168–1694.
14. Nesto R. CHD: a major burden in type 2 diabetes. *Acta Diabetol.* 2001; 38: 3–8.
15. NRC (National Research Council) Institute of Laboratory Animal Resources, Commission on Life Sciences. Guide for the Care and Use of Laboratory Animals. Washington, D.C: National Academy Press; 1996.

16. Samarghandian S, Hadjzadeh M, Amin Nya F, Davoodi S. Antihyperglycemic and antihyperlipidemic effects of guar gum on streptozotocin-induced diabetes in male rats. *Pharmacogn Mag.* 2012; 8: 65-72.